

TWO ESTIMATES FOR INTEGRAL MEANS OF  
ANALYTIC FUNCTIONS IN THE UNIT DISC

Stojan Duborija

**Abstract.** In this note two estimates for integral means of analytic functions in the unit disc are obtained. It is also shown that both estimates are sharp.

If  $f$  is an analytic function in the unit disc  $D = \{z : |z| < 1\}$ , as usual, integral mean of order  $p$ ,  $0 < p < \infty$ , of function  $f$ , is defined by

$$M_p(r, f) = \left( \frac{1}{2\pi} \int_0^{2\pi} |f(re^{i\theta})|^p d\theta \right)^{1/p}, \quad 0 < r < 1.$$

To prove our main result we need two Littlewood-Paley's theorems.

**THEOREM 1.** [2, p. 315] *Let  $0 < p \leq 2$ . Then there exists a constant  $C > 0$  such that*

$$M_p^p(r, f) \leq C \int_0^1 (1 - \rho)^{p-1} M_p^p(r\rho, f') d\rho, \quad (1)$$

for every  $f$  analytic in  $D$ .

**THEOREM 2.** [2, p. 316] *Let  $2 \leq p < \infty$ . There exists a constant  $C > 0$  such that*

$$M_p^2(r, f) \leq C \int_0^1 (1 - \rho) M_p^2(r\rho, f') d\rho, \quad (2)$$

for every  $f$  analytic in  $D$ .

The following Lemma will also be needed.

**LEMMA 1.** [2] *If  $0 < p < \infty$ , then there exists a constant  $C > 0$  such that*

$$\int_0^1 (1 - \rho)^{p-1} (1 - r\rho)^{-p} d\rho \leq C \log \frac{1}{1-r}, \quad 0 < r < 1.$$

THEOREM. Let  $M_p(r, f') = O((1-r)^{-1})$ . Then

a) if  $0 < p \leq 2$ , then  $M_p^p(r, f) = O(\log(1-r)^{-1})$ ;

b) if  $2 \leq p < \infty$ , then  $M_p^2(r, f) = O(\log(1-r)^{-1})$ .

Both estimates are sharp.

Note that from the same assumption  $M_p(r, f') = O((1-r)^{-1})$  two different conclusions follows:

$$M_p(r, f) = O\left(\log^{1/p} \frac{1}{1-r}\right), \quad \text{if } 0 < p \leq 2,$$

$$M_p(r, f) = O\left(\log^{1/2} \frac{1}{1-r}\right), \quad \text{if } 2 \leq p < \infty.$$

*Proof.* a) Using (1) we find that

$$M_p^p(r, f) \leq C \int_0^1 (1-\rho)^{p-1} (1-r\rho)^{-p} d\rho.$$

Now a) follows by Lemma 1.

We use  $C$  to denote a positive constant not necessarily the same on any two occurrences.

b) By assumption,

$$M_p(r\rho, f') \leq \frac{C}{1-r\rho}. \quad (3)$$

From (2) and (3) we conclude that

$$M_p^2(r, f) \leq C \int_0^1 (1-\rho)(1-r\rho)^{-2} d\rho.$$

Using Lemma 1 ( $p = 2$ ) and (3) we find that  $M_p^2(r, f) \leq C \log((1-r)^{-1})$ .

We show that both estimates are sharp in the sense that there exists  $f$ , analytic in  $D$ , such that  $M_p(r, f') \leq C(1-r)^{-1}$ ,  $0 < p \leq 2$ , and

$$C^{-1} \log \frac{1}{1-r} \leq M_p^p(r, f) \leq C \log \frac{1}{1-r}, \quad (4)$$

and that there exists  $g$ , analytic in  $D$ , such that  $M_p(r, g') \leq C(1-r)^{-1}$ ,  $2 \leq p < \infty$ , and

$$C^{-1} \log \frac{1}{1-r} \leq M_p^2(r, f) \leq C \log \frac{1}{1-r}. \quad (5)$$

We show that the functions  $f$  and  $g$  defined by  $f(z) = (1-z)^{-1/p}$  and  $g(z) = \sum_{n=0}^{\infty} z^{2^n}$  satisfy conditions mentioned above. Using Lemma [1, p. 65] we find that

$$M_p(r, f') \leq C \left( \frac{1}{2\pi} \int_0^{2\pi} \frac{dt}{|1 - re^{it}|^{p+1}} \right)^{1/p} \leq \frac{C}{1-r}.$$

On the other hand,  $M_p^p(r, f) = (2\pi)^{-1} \int_0^{2\pi} dt/|1 - re^{it}|$ . It is easily seen that there exists a constant  $C > 0$  so that

$$C^{-1} \log \frac{1}{1-r} \leq \int_0^{2\pi} \frac{dt}{|1 - re^{it}|} \leq C \log \frac{1}{1-r}.$$

Hence, we obtain (4).

Using inequality  $M_p(r, g') \leq M_\infty(r, g')$ , where  $M_\infty(r, g') = \sup_{0 \leq t \leq 2\pi} |g'(re^{it})|$ , we find that

$$M_p(r, g') \leq \sum_{n=0}^{\infty} 2^n r^{2^n - 1} \leq C \frac{1}{1-r}$$

(for the last inequality see [1, p. 66]). In [2, p. 316] it is proved that (5) is valid.

This completes the proof of the Theorem. ■

#### REFERENCES

- [1] P. Duran, *Theory of  $H^p$  spaces*, Academic Press, New York 1970.
- [2] A. Zygmund, *Trigonometric Series*, Volume III, University Press, Cambridge 1959.

(received 07.04.1989, in revised form 23.12.1994.)

University of Montenegro, Faculty of Mathematical and Physical Science, 81000 Podgorica,  
P.O.Box 211, Yugoslavia